

Modular anomaly equations and S-duality in $\mathcal{N} = 2$ conformal SQCD

S. K. Ashok¹, M. Billó², E. Dell'Aquila¹, M. Frau², A. Lerda^{3,2}, M. Raman¹

¹ *Institute of Mathematical Sciences
C. I. T. Campus, Taramani
Chennai, India 600113*

² *Università di Torino, Dipartimento di Fisica
and I.N.F.N. - sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy*

³ *Università del Piemonte Orientale, Dipartimento di Scienze e Innovazione Tecnologica,
and I.N.F.N. - Gruppo Collegato di Alessandria - sezione di Torino
Viale T. Michel 11, I-15121 Alessandria, Italy*

sashok,edellaquila,madhur@imsc.res.in; billo,frau,lerda@to.infn.it

ABSTRACT: We use localization techniques to study the non-perturbative properties of an $\mathcal{N} = 2$ superconformal gauge theory with gauge group $SU(3)$ and six fundamental flavours. The instanton corrections to the prepotential, the dual periods and the period matrix are calculated in a locus of special vacua possessing a \mathbb{Z}_3 symmetry. In a semi-classical expansion, we show that these observables are constrained by S-duality via a modular anomaly equation which takes the form of a recursion relation. The solutions of the recursion relation are quasi-modular functions of $\Gamma_1(3)$, which is a subgroup of the S -duality group and is also a congruence subgroup of $SL(2, \mathbb{Z})$.

KEYWORDS: $\mathcal{N} = 2$ SYM theories, recursion relations, instantons, S-duality.

Contents

1. Introduction	1
2. The $\mathcal{N} = 2$ SU(3) theory with $N_f = 6$	3
3. The Seiberg-Witten curve	7
3.1 Derivation from M-theory	7
3.2 Relating the UV and IR parameters	10
4. Electric-magnetic duality	11
4.1 The S-duality group	12
4.1.1 An important subgroup: $\Gamma_1(3)$	15
5. The dual periods and S-duality	15
5.1 S-duality and modularity	17
5.1.1 The S' transformation	17
5.1.2 The S transformation	18
5.2 Recursion relation	19
6. Summary and discussion	21
A. Observables in the special vacuum	22
B. Modular forms of $\Gamma_1(3)$	25
C. The coefficients g_3 and g_4	28

1. Introduction

Recently there has been a lot of progress in resumming instanton effects in conformal $\mathcal{N} = 2$ supersymmetric gauge theories.¹ Most of this work has been done in the SU(2) theory with four fundamental flavours [2] - [5] and in the SU(N) theories with an adjoint matter hypermultiplet, also called $\mathcal{N} = 2^*$ theories [6] - [8]. In these analyses, the effective prepotential is organized as an expansion in inverse powers of the vacuum expectation values of the scalar field in the gauge vector multiplet and the non-perturbative contributions are computed using localization methods à la Nekrasov [9] - [12]. The resulting expressions are functions of the ultraviolet (UV) gauge coupling τ_0 or, equivalently, of the instanton counting parameter $q_0 = e^{2\pi i \tau_0}$. Quite remarkably, it has been shown that these functions,

¹For a recent review on the exact results obtained in $\mathcal{N} = 2$ theories we refer to [1] and references therein.

after resumming the instanton corrections, become quasi-modular forms of the effective infrared (IR) coupling of the massless theory on which the modular group $\Gamma = \text{SL}(2, \mathbb{Z})$ faithfully acts in the standard fashion.

For $\mathcal{N} = 2$ conformal SQCD theories with gauge groups of higher rank, progress along these lines has been difficult because the S-duality group is no longer the full modular group Γ but a discrete subgroup of $\text{SL}(2, \mathbb{R})$. In this paper, we study the simplest of such theories, namely the $\mathcal{N} = 2$ super Yang-Mills theory with gauge group $\text{SU}(3)$ and six fundamental flavours, which we describe in Section 2. At a generic point on the Coulomb branch of its moduli space, one expects that the effective IR dynamics of the theory is encoded by a Seiberg-Witten (SW) curve [13, 14] which, for the case at hand, is a surface of genus 2. We engineer such a curve in Section 3 using the NS5 – D4 brane set-up in Type IIA string theory and its up-lift to M-theory [15]. The SW curve obtained in this way depends on the masses of the matter hypermultiplets, on the Coulomb branch parameters, and on the bare coupling constant τ_0 through the parameter q_0 .

It is worth recalling that in the literature there have been several different proposals [16]–[19] for SW curves for superconformal theories, like the $\text{SU}(3)$ model we are considering. These curves are always written in terms of an effective parameter called τ . Although these various proposals appear to be quite different, in [20] it was shown that it is possible to map them onto each other by means of non-perturbative reparametrizations of the τ parameter, usually accompanied by a redefinition of the masses and of the vacuum expectation values in the Coulomb branch. However, these reparametrizations in general do not preserve global identifications on the τ space; in other words, the S-duality group one finds depends crucially on the choice of τ [21].

In this paper, in order to exhibit the duality properties in a transparent manner, we choose as τ the effective coupling constant of the massless theory in the so-called special vacuum [20]. This is a locus on the Coulomb branch that possesses a \mathbb{Z}_3 symmetry and, in terms of the classical vacuum expectation values A_u of the adjoint scalar in the $\text{U}(3)$ theory, it corresponds to setting

$$A_u = \omega^{u-1}a \quad \text{for } u = 1, 2, 3, \quad (1.1)$$

where a is an arbitrary scale and ω is a cube root of unity. An added feature of the special vacuum is that, after decoupling a $\text{U}(1)$ factor and reducing the gauge symmetry to $\text{SU}(3)$, the period matrix Ω of the massless theory turns out to be proportional to the $\text{SU}(3)$ Cartan matrix even when perturbative and instanton corrections are taken into account. Namely

$$\Omega = \tau \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}. \quad (1.2)$$

In the classical theory the prefactor is simply the bare gauge coupling τ_0 , but in the effective theory it receives a perturbative correction at 1-loop and an infinite series of non-perturbative corrections due to instantons which can be explicitly computed using localization methods. Adding them up, one obtains the effective massless coupling τ on which the S-duality group acts faithfully, even when the masses for the hypermultiplets are

turned on. It is interesting to note that the relation between τ_0 and τ can be recovered by comparing the M-theory curve of Section 3 and the SW curve derived in [18, 19]. Within the framework of [20], this turns out to be a particularly convenient choice of coordinates on the coupling constant space, in that it becomes possible to make contact with the results of the multi-instanton calculations in a straightforward and direct manner.

Once the appropriate SW curve is singled out, we proceed to derive the action of S-duality on the period integrals of the curve. This action was already obtained in [21], but for the sake of completeness we briefly revisit this derivation in Section 4. A key feature of the duality transformations is that they leave the Cartan form (1.2) of the period matrix invariant while acting on the τ -parameter as

$$S \quad : \quad \tau \rightarrow -\frac{1}{3\tau} \quad \text{and} \quad T \quad : \quad \tau \rightarrow \tau + 1 . \quad (1.3)$$

The duality group generated by S and T admits as a subgroup $\Gamma_1(3)$, which is also one of the congruence subgroups of Γ [22, 23]. One might therefore hope that there are observables that inherit quasi-modular properties under this subgroup and that it may be possible to resum the instanton contributions into quasi-modular forms, as was done for the $SU(2)$ case in [4, 3]. We explicitly realize this scenario in Section 5 where we show that S-duality indeed imposes strong constraints on the modular structure of several calculable quantities. In particular we find that the quantum part of the dual periods can be written in terms of quasi-modular forms of $\Gamma_1(3)$ and that symmetry requirements from S-duality considerations imply a modular anomaly equation that takes the form of a recursion relation for the coefficients of the dual periods in the large- a expansion. This recursion relation can then be used to calculate the higher order coefficients and verify that their instanton expansion completely agrees with direct computations performed using equivariant localization methods. We regard this agreement as a highly non-trivial consistency check of our results.

In Section 6 we present our conclusions and discuss some future perspectives, while several technical details are collected in the appendices.

2. The $\mathcal{N} = 2$ $SU(3)$ theory with $N_f = 6$

We begin by reviewing the main features of the $\mathcal{N} = 2$ superconformal theory with gauge group $U(3)$ and six fundamental flavours.

As usual it is convenient to combine the Yang-Mills coupling g and the vacuum angle θ into the complex variable

$$\tau_0 = \frac{\theta}{2\pi} + i \frac{4\pi}{g^2} , \quad (2.1)$$

and define the instanton counting parameter

$$q_0 = e^{2\pi i \tau_0} . \quad (2.2)$$

The prepotential of the theory is the sum of a classical term F_{class} , a perturbative 1-loop term $F_{1\text{-loop}}$ and a non-perturbative piece F_{inst} :

$$F = F_{\text{class}} + F_{1\text{-loop}} + F_{\text{inst}} . \quad (2.3)$$

The classical prepotential is

$$F_{\text{class}} = \pi i \tau_0 \sum_{u=1}^3 A_u^2 \quad (2.4)$$

with A_u being the vacuum expectation value of the scalar field A in the adjoint representation of $U(3)$:

$$\langle A \rangle = \text{diag}(A_1, A_2, A_3) . \quad (2.5)$$

The 1-loop term is independent of τ_0 and is given by

$$F_{1\text{-loop}} = \sum_{u \neq v} \gamma(A_u - A_v) - \sum_{u=1}^3 \sum_{f=1}^6 \gamma(A_u + m_f) , \quad (2.6)$$

where

$$\gamma(x) = -\frac{x^2}{4} \log\left(\frac{x^2}{\Lambda^2}\right) , \quad (2.7)$$

with Λ being an arbitrary mass scale. Expanding $F_{1\text{-loop}}$ for small masses, one obtains an expression in which the three vacuum expectation values A_u appear through the Casimirs of the $U(3)$ gauge group

$$u_r = \sum_{u=1}^3 A_u^r \quad \text{for } r \in \{1, \dots, 3\} , \quad (2.8)$$

and the six fundamental masses m_f through the Casimirs of the $U(6)$ flavour symmetry group

$$T_\ell = \sum_{f=1}^6 m_f^\ell \quad \text{for } \ell \in \{1, \dots, 6\} . \quad (2.9)$$

For simplicity in this paper we always consider mass configurations such that

$$T_1 = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 = 0 . \quad (2.10)$$

Thus, the flavour symmetry group is reduced to $SU(6)$.

The non-perturbative instanton part is computed using equivariant localization methods [9] - [12] (see also [24] for technical details), and the result is of the form

$$F_{\text{inst}} = \sum_{k=1}^{\infty} q_0^k f_k(u_r, T_\ell) , \quad (2.11)$$

where the coefficients f_k are rational functions of mass dimension 2 in which the Casimirs T_ℓ appear only in the numerator. We have computed these functions up to $k = 4$, but we do not write them here since their explicit expressions are too lengthy².

If the gauge group is $SU(3)$ instead of $U(3)$, we simply have to enforce the constraint

$$u_1 = A_1 + A_2 + A_3 = 0 \quad (2.12)$$

²However, they can be made available upon request.

in all above formulas. We do this by setting

$$A_1 = a_1 , \quad A_2 = a_2 - a_1 , \quad A_3 = -a_2 , \quad (2.13)$$

which corresponds to choosing the two Cartan generators of $SU(3)$ as $H_1 = \text{diag}(1, -1, 0)$ and $H_2 = \text{diag}(0, 1, -1)$. With this choice the classical prepotential (2.4) becomes

$$F_{\text{class}} = 2\pi i \tau_0 (a_1^2 - a_1 a_2 + a_2^2) , \quad (2.14)$$

from which we easily see that the classical period matrix is

$$(\Omega_{\text{class}})_{ij} := \frac{1}{2\pi i} \frac{\partial^2 F_{\text{class}}}{\partial a_i \partial a_j} = \tau_0 \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}_{ij} , \quad (2.15)$$

namely Ω_{class} is proportional to the Cartan matrix of $SU(3)$.

When we add the perturbative and instanton corrections to the classical prepotential, this structure is generically lost. However there is a locus in the moduli space where the period matrix retains a simple form: it is the so-called *special vacuum* [20] which corresponds to setting

$$u_2 = 0 . \quad (2.16)$$

This condition can be achieved by taking a \mathbb{Z}_3 -symmetric configuration of the A_u 's in the original $U(3)$ theory, namely

$$A_u = \omega^{u-1} a , \quad (2.17)$$

where $\omega = e^{\frac{2\pi i}{3}}$ and a is an arbitrary scale. In terms of the $SU(3)$ variables introduced in (2.13), the special vacuum corresponds to

$$a_1 = a , \quad a_2 = -\omega^2 a . \quad (2.18)$$

With this choice, the classical prepotential (2.14) vanishes, while the 1-loop term is³

$$F_{1\text{-loop}} \Big|_{\text{s.v.}} = \frac{T_2}{2} \log \left(\frac{a^3}{\Lambda^3} \right) + \frac{T_5}{20a^3} - \frac{T_8}{112a^6} + \frac{T_{11}}{330a^9} + \dots \quad (2.19)$$

where the T_ℓ 's with $\ell > 6$ are defined as in (2.9) and can be written as polynomials of the independent lower Casimirs. Adding the instanton corrections, the full prepotential in the special vacuum becomes

$$F \Big|_{\text{s.v.}} = \frac{T_2}{2} \log \left(\frac{a^3}{\Lambda^3} \right) + \sum_{n=1}^{\infty} \frac{\hat{h}_n(q_0, T_\ell)}{n a^{3n}} \quad (2.20)$$

where the first few coefficients \hat{h}_n are reported, up to four instantons, in Appendix A (see in particular (A.2)). The period matrix Ω in the special vacuum takes a rather simple form; defining the matrices

$$\mathcal{C} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} , \quad \mathcal{B} = \begin{pmatrix} \omega^2 & \omega \\ \omega & 1 \end{pmatrix} \quad \text{and} \quad \mathcal{B}^\dagger = \begin{pmatrix} \omega & \omega^2 \\ \omega^2 & 1 \end{pmatrix} , \quad (2.21)$$

³Here and in the following we always neglect all a -independent terms.

one finds:

$$\Omega = \frac{1}{2\pi i} \left(2\pi i \tau_0 - i\pi - \log 27 + \frac{4q_0}{9} + \frac{14q_0^2}{81} + \frac{1948q_0^3}{19683} + \dots \right) \mathcal{C} \quad (2.22)$$

$$- \frac{1}{2\pi i} \sum_{n=0}^{\infty} \left[\left(\frac{3n+1}{a^{3n+2}} \hat{h}_n(q_0, T_\ell) \right) \mathcal{B} + \left(\frac{3n+2}{a^{3n+3}} \hat{g}_n(q_0, T_\ell) \right) \mathcal{C} + \left(\frac{3n+3}{a^{3n+4}} \hat{k}_n(q_0, T_\ell) \right) \mathcal{B}^\dagger \right].$$

The $\hat{h}_n(q_0, T_\ell)$'s are the same functions that appear in the prepotential (2.20), while $\hat{g}_n(q_0, T_\ell)$ and $\hat{k}_n(q_0, T_\ell)$ can be reconstructed from the expressions given in Appendix A (see (A.5) and (A.7)). Looking at them, it is easy to realize that the coefficients of \mathcal{B} and \mathcal{B}^\dagger vanish if $T_2 = T_4 = T_5 = 0$.

Prompted by this observation, we conclude that turning on only T_3 and T_6 yields a period matrix proportional to \mathcal{C} with the constant of proportionality playing the role of an effective IR coupling constant, even after taking into account the 1-loop and instanton contributions. In terms of the bare masses, this is equivalent to making a \mathbb{Z}_3 -symmetric choice, for instance

$$(m_1, m_2, m_3) = (m, m\omega, m\omega^2) \quad \text{and} \quad (m_4, m_5, m_6) = (m', m'\omega, m'\omega^2), \quad (2.23)$$

for which the non-zero Casimirs T_3 and T_6 become

$$T_3 = 3(m^3 + m'^3) \quad \text{and} \quad T_6 = 3(m^6 + m'^6). \quad (2.24)$$

With this mass configuration, the period matrix maintains the same simple form (2.15) of the classical theory, namely

$$\Omega = \tilde{\tau} \mathcal{C}, \quad (2.25)$$

where the effective IR coupling $\tilde{\tau}$ is

$$\begin{aligned} 2\pi i \tilde{\tau} = & 2\pi i \tau_0 - i\pi - \log 27 + \frac{4q_0}{9} + \frac{14q_0^2}{81} + \frac{1948q_0^3}{19683} + \dots \\ & + \frac{T_3}{a^3} \left(\frac{1}{3} - \frac{2q_0}{27} - \frac{8q_0^2}{243} - \frac{398q_0^3}{19683} + \dots \right) - \frac{T_6}{a^6} \left(\frac{1}{6} + \frac{5q_0}{27} + \frac{5q_0^2}{243} + \frac{5q_0^3}{19683} + \dots \right) \\ & + \frac{T_3^2}{a^6} \left(\frac{5q_0}{81} + \frac{10q_0^2}{729} + \frac{95q_0^3}{19683} + \dots \right) + \dots \end{aligned} \quad (2.26)$$

A quantity of interest, which will play a crucial role in later sections, is the effective IR coupling of the *massless* theory. Notice that even when all masses are zero there is a non-perturbative renormalization of the bare UV coupling τ_0 to an effective IR coupling⁴. We will denote this coupling by τ which, as we can see from (2.26) with $T_3 = T_6 = 0$, is given by

$$2\pi i \tau = 2\pi i \tau_0 - i\pi - \log 27 + \frac{4q_0}{9} + \frac{14q_0^2}{81} + \frac{1948q_0^3}{19683} + \dots \quad (2.27)$$

⁴This is different from the $\mathcal{N} = 2^*$ theories [3, 4, 6] where for zero mass one recovers the maximal $\mathcal{N} = 4$ supersymmetry which implies that the coupling constant does not run, neither perturbatively nor non-perturbatively.

This relation can be inverted and, exponentiating both sides we find

$$q_0 = -27 q (1 + 12q + 90q^2 + 508q^3 + \dots) , \quad (2.28)$$

where $q = e^{2\pi i \tau}$. As we shall see in the following sections, the massless IR coupling τ plays an important role because it is the natural parameter that transforms faithfully under the S-duality transformations of the effective theory [25].

3. The Seiberg-Witten curve

In this section we derive the SW curve for the $SU(3)$ gauge theory with $N_f = 6$ flavours discussed before. We engineer such a theory using the NS5 – D4 brane set up of Type IIA string theory [15], and closely follow the notations and conventions of [26].

3.1 Derivation from M-theory

Our construction begins with a collection of NS5 and D4 branes, arranged as indicated in Tab. 1. The first four directions $\{x^0, x^1, x^2, x^3\}$ are longitudinal for both kinds of branes

	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9	x^{10}
NS5 branes	—	—	—	—	—	—	·	·	·	·	·
D4 branes	—	—	—	—	·	·	—	·	·	·	—

Table 1: Type IIA brane configuration: the symbols — and · denote longitudinal and transverse directions respectively; the last column refers to the eleventh dimension after the M-theory uplift.

and span the spacetime $\mathbb{R}^{1,3}$ where the gauge theory is defined. After compactifying the x^5 direction on a circle S^1 of radius R_5 , we uplift the system to M-theory by introducing a compact eleventh coordinate x^{10} with radius R_{10} . We then minimize the world-volume of the resulting M5 branes and in doing so, obtain the SW curve for a 5-dimensional $\mathcal{N} = 1$ gauge theory defined in $\mathbb{R}^{1,3} \times S^1$ which takes the form of a 2-dimensional surface inside the space parameterized by the coordinates $\{x^4, x^5, x^6, x^{10}\}$. To get the curve for the 4-dimensional $\mathcal{N} = 2$ theory, we first T-dualize along x^5 and then take the limit of small (dual) radius. Thus, in terms of the dual circumference

$$\beta = \frac{2\pi\alpha'}{R_5} , \quad (3.1)$$

the 4-dimensional limit corresponds to $\beta \rightarrow 0$.

Let us now give some details. To engineer the conformal $SU(3)$ gauge theory, we consider the following configuration (see Fig. 1):

- Two NS5 branes separated by finite distances along the x^6 direction, labeled NS5₁ and NS5₂,
- Three semi-infinite D4 branes on NS5₁ and three semi-infinite D4 branes on the NS5₂,




Figure 1: NS5 and D4 brane set up for the conformal SU(3) theory.

- Three finite D4 branes suspended between NS5₁ and NS5₂.

The brane configuration is best described in terms of the complex combinations

$$x^4 + i x^5 \equiv 2\pi\alpha'v \quad \text{and} \quad x^6 + i x^{10} \equiv s, \quad (3.2)$$

or their exponentials

$$w \equiv e^{\frac{2\pi\alpha'v}{R_5}} = e^{\beta v} \quad \text{and} \quad t \equiv e^{\frac{s}{R_{10}}}, \quad (3.3)$$

which are single-valued under integer shifts of x^5 and x^{10} along the respective circumferences.

The SW curve for the T-dual 5-dimensional gauge theory takes the form of a polynomial equation of degree 2 in t , since there are two NS5 branes, and of degree 3 in w , since there are three suspended D4 branes. Such a polynomial equation can be written in two distinct ways; the first is [26]

$$\mathcal{C}_1 : \quad w^3 (t - t_1^{(\infty)}) (t - t_2^{(\infty)}) + w^2 Q_2(t) + w Q_1(t) + d' (t - t_1^{(0)}) (t - t_2^{(0)}) = 0. \quad (3.4)$$

Here, $Q_1(t)$ and $Q_2(t)$ are quadratic polynomials in t , d' is an arbitrary coefficient, while $t_i^{(0)}$ and $t_i^{(\infty)}$ are the asymptotic positions of the two NS5 branes as $w \rightarrow 0$ and $w \rightarrow \infty$

respectively. These are given by

$$\begin{aligned} t_1^{(\infty)} &= C^{(1)} \sqrt{\frac{\tilde{m}_1 \tilde{m}_2 \tilde{m}_3}{\tilde{A}_1 \tilde{A}_2 \tilde{A}_3}}, & t_1^{(0)} &= C^{(1)} \sqrt{\frac{\tilde{A}_1 \tilde{A}_2 \tilde{A}_3}{\tilde{m}_1 \tilde{m}_2 \tilde{m}_3}}, \\ t_2^{(\infty)} &= C^{(2)} \sqrt{\frac{\tilde{A}_1 \tilde{A}_2 \tilde{A}_3}{\tilde{m}_4 \tilde{m}_5 \tilde{m}_6}}, & t_2^{(0)} &= C^{(2)} \sqrt{\frac{\tilde{m}_4 \tilde{m}_5 \tilde{m}_6}{\tilde{A}_1 \tilde{A}_2 \tilde{A}_3}} \end{aligned} \quad (3.5)$$

where $C^{(1)}$ and $C^{(2)}$ are constants and the tilded variables are defined as

$$\tilde{X} = e^{\beta X} \quad (3.6)$$

for any X . In particular the tilded variables of (3.5) correspond to the positions of the D4 branes in the (x^4, x^5) -plane which represent the three vacuum expectation values A_u of the adjoint scalar field and the six masses m_f of the U(3) gauge theory, as described in the previous section. Since we are interested in the SU(3) theory, we impose the constraint (2.12) which corresponds to setting the center of mass of the colour D4 branes to the origin.

The second form of the SW curve is [26]

$$\mathcal{C}_2 : (w - \tilde{m}_1)(w - \tilde{m}_2)(w - \tilde{m}_3)t^2 + P(w)t + d(w - \tilde{m}_4)(w - \tilde{m}_5)(w - \tilde{m}_6) = 0, \quad (3.7)$$

where $P(w)$ is a cubic polynomial and d is an arbitrary coefficient. This structure follows from the fact that when $t \rightarrow 0$ and $t \rightarrow \infty$ there are three D4 flavour branes at $w = (\tilde{m}_1, \tilde{m}_2, \tilde{m}_3)$ and $w = (\tilde{m}_4, \tilde{m}_5, \tilde{m}_6)$, respectively.

Equating (3.5) and (3.7), we can fix all but two coefficients in terms of the physical parameters of the gauge theory and get

$$\begin{aligned} &(w - \tilde{m}_1)(w - \tilde{m}_2)(w - \tilde{m}_3)t^2 \\ &- \left[(t_1^{(\infty)} + t_2^{(\infty)})w^3 + c_2 w^2 + c_1 w - \tilde{m}_1 \tilde{m}_2 \tilde{m}_3 (t_1^{(0)} + t_2^{(0)}) \right] t \\ &+ t_1^{(\infty)} t_2^{(\infty)} (w - \tilde{m}_4)(w - \tilde{m}_5)(w - \tilde{m}_6) = 0 \end{aligned} \quad (3.8)$$

where the two undetermined coefficients, c_1 and c_2 , parametrize the Coulomb branch of the SU(3) gauge theory. According to [15], the difference in asymptotic positions of the NS5 branes is related to the bare gauge coupling constant; to be precise, one has

$$q_0 = \frac{C^{(1)}}{C^{(2)}}. \quad (3.9)$$

Now we can take the four dimensional limit $\beta \rightarrow 0$. Skipping several intermediate steps which are similar to those described in [26], we find that the four dimensional SW curve for the SU(3) superconformal theory is

$$(v - m_1)(v - m_2)(v - m_3)t^2 - ((1 + q_0)v^3 - U_2 v - U_3)t + q_0(v - m_4)(v - m_5)(v - m_6) = 0. \quad (3.10)$$

Here we have used the fact that $T_1 = \sum_f m_f = 0$ and have redefined the coefficients c_1 and c_2 to absorb all terms linear in t and proportional to v or independent of v into the new parameters U_2 and U_3 . These have been normalized in such a way that

$$U_2|_{q_0 \rightarrow 0} = u_2 \quad \text{and} \quad U_3|_{q_0 \rightarrow 0} = u_3 . \quad (3.11)$$

In the special vacuum and in the \mathbb{Z}_3 -symmetric mass configuration (2.23), the SW curve (3.10) further simplifies to

$$(v^3 - m^3) t^2 - ((1 + q_0)v^3 - U_3) t + q_0(v^3 - m'^3) = 0 . \quad (3.12)$$

3.2 Relating the UV and IR parameters

The SW curve (3.10) is written in terms of the UV parameters of the $SU(3)$ gauge theory; in particular it explicitly depends on q_0 which is the instanton counting parameter used in the Nekrasov analysis of the non-perturbative prepotential. However, it would be useful to have a description in terms of the effective IR parameters and couplings since these are the ones which faithfully transform under the duality group of the conformal gauge theory.

To do this let us first rewrite the curve (3.10) in a different form [27]. Define the monic polynomial

$$\mathcal{P}(v) = v^3 - \frac{U_2}{1 + q_0} v - \frac{U_3}{1 + q_0} \quad (3.13)$$

and rescale t into $(1 + q_0)t$, so that (3.10) becomes

$$(v - m_1)(v - m_2)(v - m_3) t^2 - \mathcal{P}(v) t + \frac{q_0}{(1 + q_0)^2} (v - m_4)(v - m_5)(v - m_6) = 0 . \quad (3.14)$$

If we shift away the term linear in t and introduce

$$y = 2(v - m_1)(v - m_2)(v - m_3) \left(t - \frac{\mathcal{P}(v)}{2(v - m_1)(v - m_2)(v - m_3)} \right) , \quad (3.15)$$

we obtain

$$y^2 = \mathcal{P}^2(v) - \frac{4q_0}{(1 + q_0)^2} \mathcal{Q}(v) \quad (3.16)$$

where

$$\mathcal{Q}(v) = \prod_{f=1}^6 (v - m_f) . \quad (3.17)$$

In this way we manage to cast the M-theory curve for our superconformal $SU(3)$ gauge theory into a hyperelliptic form which makes comparison with the earlier literature [16, 17, 18, 19] straightforward. In particular, consider the curve given in Eq. (4.12) of [19]. This curve is written in terms of the IR coupling τ of the massless theory in the special vacuum and, after a shift and rescaling, it takes the form

$$y^2 = \mathcal{P}^2(v) - h(\tau) \mathcal{Q}(v) \quad (3.18)$$

where

$$h(\tau) = \left(\frac{f_-^2(\tau) - f_+^2(\tau)}{f_-^2(\tau)} \right) . \quad (3.19)$$

Here $f_{\pm}(\tau)$ are modular forms of $\Gamma_1(3)$ with weight 3, given by

$$f_{\pm}(\tau) = \left(\frac{\eta^3(\tau)}{\eta(3\tau)} \right)^3 \pm \left(\frac{3\eta^3(3\tau)}{\eta(\tau)} \right)^3 \quad (3.20)$$

with $\eta(\tau)$ being the Dedekind η -function [19] (see also [22, 23]). We refer to Appendix B for the properties of the modular functions and a brief review of $\Gamma_1(3)$ which, as we show later, arises as a subgroup of the S-duality group of the $SU(3)$ superconformal theory.

The two curves (3.16) and (3.18) become identical if

$$\frac{f_+(\tau)}{f_-(\tau)} = \frac{1 - q_0}{1 + q_0} , \quad (3.21)$$

which implies

$$q_0 = -27 \left(\frac{\eta(3\tau)}{\eta(\tau)} \right)^{12} . \quad (3.22)$$

This relation has already been obtained in [24] using different arguments related to the holographic description of the effective gauge coupling. Fourier expanding the η -functions in powers of $q = e^{2\pi i \tau}$, we obtain

$$q_0 = -27 q (1 + 12q + 90q^2 + 508q^3 + \dots) \quad (3.23)$$

which exactly agrees with (2.28). The fact that the parameter τ appearing in the modular functions of the curve (3.18) is related to the effective IR coupling of the massless theory in the special vacuum (and hence to the period matrix Ω) provides a nice geometrical interpretation, while the fact that the UV/IR relation (3.22) is in perfect agreement with the explicit multi-instanton calculations gives a strong confirmation on the validity of the whole picture.

4. Electric-magnetic duality

The SW curve (3.18) represents a genus 2 surface on which we can choose a canonical homology basis of cycles $\{\alpha_i\}$ and $\{\beta_j\}$ (with $i, j = 1, 2$) such that $\alpha_i \cap \beta_j = \delta_{ij}$. On these cycles the $Sp(4, \mathbb{Z})$ electric-magnetic duality group acts in the standard fashion. We denote an element of this group by

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(4, \mathbb{Z}) , \quad (4.1)$$

where A, B, C and D are 2×2 matrices satisfying the relations

$$A^T C - C^T A = 0 , \quad B^T D - D^T B = 0 , \quad \text{and} \quad A^T D - C^T B = \mathbb{1}_2 . \quad (4.2)$$

Under $Sp(4, \mathbb{Z})$ the homology cycles transform as

$$\begin{pmatrix} \beta \\ \alpha \end{pmatrix} \rightarrow M \cdot \begin{pmatrix} \beta \\ \alpha \end{pmatrix} = \begin{pmatrix} A\beta + B\alpha \\ C\beta + D\alpha \end{pmatrix} , \quad (4.3)$$

while the period matrix transforms as

$$\Omega \longrightarrow (A\Omega + B)(C\Omega + D)^{-1}. \quad (4.4)$$

Given the SW differential λ , the period integrals over the α_i - and β_j -cycles are denoted as

$$a_i = \oint_{\alpha_i} \lambda \quad \text{and} \quad a_j^D = \oint_{\beta_j} \lambda. \quad (4.5)$$

The α -periods represent the vacuum expectation values of the adjoint scalar field; that is why we used the same notation as in Section 2. The β -periods, instead, represent their dual variables which are related to the prepotential F according to

$$a_i^D = \frac{1}{2\pi i} \frac{\partial F}{\partial a_i}. \quad (4.6)$$

From the transformation rules (4.3) of the cycles under $\text{Sp}(4, \mathbb{Z})$, it immediately follows that the period integrals transform according to

$$\begin{pmatrix} a^D \\ a \end{pmatrix} \rightarrow M \cdot \begin{pmatrix} a^D \\ a \end{pmatrix} = \begin{pmatrix} A a^D + B a \\ C a^D + D a \end{pmatrix}. \quad (4.7)$$

4.1 The S-duality group

The S-duality groups of $\mathcal{N} = 2$ conformal SQCD theories have been studied in [25, 21]. Here we will briefly review the analysis of [21] adapting it to our $\text{SU}(3)$ theory.

Referring to the SW curve (3.18), the total moduli space has the structure of a fiber bundle in which the h -plane is the base and the Coulomb branch is the fiber. In the conventional SW analysis of non-scale invariant theories [13], the monodromies around the singularities of the Coulomb branch generate the low energy electric-magnetic duality group. In scale invariant theories, like the one we are considering, there are in addition isolated “special” singularities in the space of coupling constants where the entire Coulomb branch degenerates. Monodromies around these special points in the h -plane are *defined* to be the generators of the S-duality group [21].

The discriminant locus of the SW curve (3.18) is proportional to

$$\Delta = h^3(h-1)^3. \quad (4.8)$$

Thus, the curve is singular at $h = 0, 1$ and ∞ . In the special vacuum, these are the only singularities around which the curve degenerates [21]. To discuss how S-duality is realized on our SW curve, let us temporarily return to a $\text{U}(3)$ set-up and choose a \mathbb{Z}_3 -symmetric set of cycles $\{\hat{\alpha}_u\}$ and $\{\hat{\beta}_v\}$ ($u, v = 1, 2, 3$) as shown in Fig. 2, whose intersection matrix is

$$(\hat{\alpha}_u \cap \hat{\beta}_v) = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ -1 & 0 & 1 \end{pmatrix}. \quad (4.9)$$

These cycles are not linearly independent since

$$\sum_{u=1}^3 \hat{\alpha}_u = 0 \quad \text{and} \quad \sum_{v=1}^3 \hat{\beta}_v = 0. \quad (4.10)$$




Figure 2: The $\hat{\alpha}$ and $\hat{\beta}$ cycles for the U(3) SW curve.

Under the monodromy $h \rightarrow e^{2\pi i}h$, one can check that the branch points are rotated such that the cycles are transformed as follows⁵

$$\hat{T} : \begin{cases} \hat{\alpha}_u \rightarrow \hat{\alpha}_u , \\ \hat{\beta}_v \rightarrow \hat{\beta}_v + \hat{\alpha}_v - \hat{\alpha}_{v-1} . \end{cases} \quad (4.11)$$

Under $(h-1) \rightarrow e^{2\pi i}(h-1)$, we obtain the monodromy $\hat{S}\hat{T}^{-1}$, which acts on the cycles as

$$\hat{S}\hat{T}^{-1} : \begin{cases} \hat{\alpha}_u \rightarrow \hat{\beta}_u - \hat{\alpha}_u + \hat{\alpha}_{u-1} , \\ \hat{\beta}_v \rightarrow \hat{\alpha}_{v-1} . \end{cases} \quad (4.12)$$

It then follows that the \hat{S} transformation, which corresponds to the monodromy around $h \rightarrow \infty$, is given by

$$\hat{S} : \begin{cases} \hat{\alpha}_u \rightarrow \hat{\beta}_u , \\ \hat{\beta}_v \rightarrow \hat{\alpha}_{v-1} . \end{cases} \quad (4.13)$$

We have already remarked that the Riemann surface is invariant under a \mathbb{Z}_3 rotation of the cycles generated by

$$\hat{\Phi} : \begin{cases} \hat{\alpha}_u \rightarrow \hat{\alpha}_{u-1} , \\ \hat{\beta}_v \rightarrow \hat{\beta}_{v-1} . \end{cases} \quad (4.14)$$

Therefore, instead of the transformation \hat{S} we can equivalently consider the transformation obtained by composing it with $\hat{\Phi}$, *i.e.* we define a modified \hat{S} transformation

$$\hat{S} \rightarrow \hat{\Phi} \cdot \hat{S} : \begin{cases} \hat{\alpha}_u \rightarrow \hat{\beta}_{u-1} , \\ \hat{\beta}_v \rightarrow \hat{\alpha}_{v+1} . \end{cases} \quad (4.15)$$

⁵Here we use the convention that the indices u and v are defined modulo 3 and that the hatted operators act on the U(3) basis of cycles while the unhatted ones correspond to $\text{Sp}(4, \mathbb{Z})$ matrices which are obtained by restricting the action of the hatted operators to the SU(3) basis which we introduce later.

For simplicity $\hat{\Phi} \cdot \hat{S}$ will be renamed \hat{S} . The new duality transformations satisfy

$$\hat{S}^2 = 1 \quad \text{and} \quad (\hat{S}\hat{T})^6 = 1 . \quad (4.16)$$

Let us now return to the SU(3) theory and choose the independent α - and β -cycles as

$$\begin{aligned} \alpha_1 &= \hat{\alpha}_1 & \text{and} & & \alpha_2 &= -\hat{\alpha}_3 , \\ \beta_1 &= -\hat{\beta}_2 & \text{and} & & \beta_2 &= -\hat{\beta}_3 . \end{aligned} \quad (4.17)$$

Using (4.9), it is easy to verify that $\alpha_i \cap \beta_j = \delta_{ij}$, as it should be for a canonical homology basis. The restrictions of the \hat{S} and \hat{T} transformations to the SU(3) basis (4.17) follow directly from the transformation rules (4.11) and (4.15) in the parent U(3) theory. Representing them as 4×4 matrices acting on the 4-vector $\begin{pmatrix} \beta \\ \alpha \end{pmatrix}$, we have

$$T = \begin{pmatrix} 1 & 0 & 2 & -1 \\ 0 & 1 & -1 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad S = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} . \quad (4.18)$$

Acting projectively as described in (4.4) on the period matrix $\Omega = \tau \mathcal{C}$ these matrices leave its matrix structure invariant, while the prefactor τ changes as follows

$$T : \tau \longrightarrow \tau + 1 \quad \text{and} \quad S : \tau \longrightarrow -\frac{1}{3\tau} . \quad (4.19)$$

These transformations satisfy the relations

$$S^2 = 1 \quad \text{and} \quad (ST)^6 = 1 , \quad (4.20)$$

just as their U(3) parents \hat{T}, \hat{S} in (4.16). They generate a discrete subgroup of $\text{SL}(2, \mathbb{R})$ which is the duality group of the theory.

As we observed earlier, this duality group has a natural action also on the periods a_i and their dual ones a_j^{D} which is inherited from the action of T and S on the α_i and β_j cycles. In particular under the S -transformation in (4.18), we have

$$S : \begin{pmatrix} a_1^{\text{D}} \\ a_2^{\text{D}} \\ a_1 \\ a_2 \end{pmatrix} \longrightarrow \begin{pmatrix} a_2 \\ -a_1 \\ -a_2^{\text{D}} \\ a_1^{\text{D}} \end{pmatrix} . \quad (4.21)$$

Note that under this transformation the a_i 's and the a_j^{D} are exchanged (up to signs), as expected for an S -transformation. Note also that the transformations (4.19) leave invariant the function $h(\tau)$ in (3.19) that appears in the SW curve (3.18) of the SU(3) theory, as can be checked using the modular properties of the Dedekind η -functions reported in Appendix B.

4.1.1 An important subgroup: $\Gamma_1(3)$

An important subgroup of the full duality group, which will play an central role in what follows, is the one generated by T (as in (4.18)) and S' where [25]

$$S' = S T S^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -2 & -1 & 1 & 0 \\ -1 & -2 & 0 & 1 \end{pmatrix}. \quad (4.22)$$

This S' rotates the periods into each other according to

$$S' : \begin{pmatrix} a_1^D \\ a_2^D \\ a_1 \\ a_2 \end{pmatrix} \longrightarrow \begin{pmatrix} a_1^D \\ a_2^D \\ a_1 - 2a_1^D - a_2^D \\ a_2 - a_1^D - 2a_2^D \end{pmatrix}, \quad (4.23)$$

while it acts on the coupling τ as

$$S' : \tau \longrightarrow \frac{\tau}{1 - 3\tau}. \quad (4.24)$$

T and S' generate the group $\Gamma_1(3)$, which is a congruence subgroup of the modular group $\text{SL}(2, \mathbb{Z})$:

$$\Gamma_1(3) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{Z}) : c = 0 \pmod{3}, a = d = 1 \pmod{3} \right\}. \quad (4.25)$$

As we have already mentioned, the function $h(\tau)$ appearing in the SW curve (3.18) is written in terms of modular forms of $\Gamma_1(3)$. In what follows, we will exploit this symmetry and in particular, using the above action of S-duality, we will show that the dual periods of the $\text{SU}(3)$ theory in the special vacuum can be expanded in terms of quasi-modular functions of $\Gamma_1(3)$.

5. The dual periods and S-duality

In order to study the S-duality properties of the superconformal $\text{SU}(3)$ theory, we have to calculate the dual periods a_i^D in the special vacuum. This is done by deriving the prepotential F and then setting the scalars to their vacuum expectation values of the special vacuum:

$$a_i^D = \frac{1}{2\pi i} \left. \frac{\partial F}{\partial a_i} \right|_{\text{s.v.}}. \quad (5.1)$$

In the special vacuum, there is only one independent pair of dual variables, which we choose to be (a_1, a_2^D) . This follows from the S-duality rules derived in (4.21). Thus, from now on we use the notation $a_1 \equiv a$ and $a_2^D \equiv a^D$.

We consider the classical, 1-loop and instanton contributions separately. From the prepotential (2.14) we find that the classical contribution is⁶

$$a_{\text{class}}^{\text{D}} = i\sqrt{3}\tau_0 a . \quad (5.2)$$

The 1-loop contribution to the dual period in the special vacuum is given by

$$a_{1\text{-loop}}^{\text{D}} = \frac{\sqrt{3}}{2\pi} \left[(-i\pi - \log 27)a - \frac{T_3}{6a^2} + \frac{T_6}{30a^5} + \frac{T_3(T_3^2 - 9T_6)}{1296a^8} + \dots \right] , \quad (5.3)$$

while the instanton corrections, which are a power series in q_0 , can be calculated using the techniques of equivariant localization.

These results, however, are not very illuminating. Things change significantly if we write the dual period using the massless effective coupling τ , which transforms faithfully under the S-duality group. To do so, we have use the UV/IR relation (3.23). Once this is done, we find that a^{D} takes the form

$$a^{\text{D}} = i\sqrt{3}\tau a + \frac{\sqrt{3}}{2\pi} \sum_{n=0}^{\infty} \frac{g_n(\tau, T_\ell)}{a^{3n+2}} , \quad (5.4)$$

where the first few g_n 's have the following q -expansions

$$g_0 = -\frac{T_3}{6} (1 + 6q + 6q^3 + \dots) , \quad (5.5a)$$

$$g_1 = \frac{T_3^2}{3} (q + 6q^2 + 3q^3 + \dots) + \frac{T_6}{30} (1 - 30q - 270q^2 - 570q^3 + \dots) , \quad (5.5b)$$

$$g_2 = \frac{T_3^3}{1296} (1 - 756q^2 - 336q^3 + \dots) - \frac{T_3 T_6}{144} (1 - 252q^2 + 672q^3 + \dots) , \quad (5.5c)$$

$$g_3 = -\frac{T_3^4}{7128} (1 + 1386q^2 - 11088q^3 + \dots) + \frac{T_3^2 T_6}{1188} (1 + 1386q^2 - 5808q^3 + \dots) \\ + \frac{T_6^2}{792} (1 - 1386q^2 + 528q^3 + \dots) , \quad (5.5d)$$

$$g_4 = \frac{T_3^5}{68040} (1 + 87360q^3 + \dots) - \frac{T_3^3 T_6}{27} (208q^3 + \dots) \\ - \frac{T_3 T_6^2}{1512} (1 - 17472q^3 + \dots) \quad (5.5e)$$

when only T_3 and T_6 are turned on. In what follows we will show that these coefficients g_n 's, which appear also in front of \mathcal{C} in the period matrix, are in fact quasi-modular functions of $\Gamma_1(3)$ with weight $3n + 1$.

⁶For completeness, we point out that in the special vacuum the period a_1^{D} is classically proportional to a_2^{D} . Even when the instanton corrections are taken into account, the following relation holds in the special vacuum:

$$a_1^{\text{D}}|_{\text{s.v.}} = \frac{1}{\omega} a_2^{\text{D}}|_{\text{s.v.}} .$$

5.1 S-duality and modularity

In Section 4 we have already derived how S-duality acts on the periods and their duals. In order to see the consequences of this when we are in the special vacuum, let us consider the combination

$$X = a^D - i\sqrt{3}\tau a = \frac{\sqrt{3}}{2\pi} \sum_n \frac{g_n(\tau, T_\ell)}{a^{3n+2}} \quad (5.6)$$

and study its transformation properties under both S and S' using the $\mathrm{Sp}(4, \mathbb{Z})$ matrices defined in (4.21) and (4.23) respectively.

5.1.1 The S' transformation

Under the S' -transformation, we find

$$\begin{aligned} a^D &\rightarrow a^D, \\ a &\rightarrow (a + i\sqrt{3}a^D). \end{aligned} \quad (5.7)$$

From this it follows that X transforms as

$$S'(X) = \frac{X}{(1 - 3\tau)}. \quad (5.8)$$

If we act with S' on the right hand side of (5.6), we find

$$S'(X) = \frac{\sqrt{3}}{2\pi} \sum_n \frac{S'(g_n(\tau, T_\ell))}{a^{3n+2}(1 - 3\tau)^{3n+2}} \left(1 - \frac{3}{2\pi i(1 - 3\tau)} \sum_m \frac{g_m}{a^{3m+3}} \right)^{-3n-2}. \quad (5.9)$$

Equating (5.8) and (5.9), we deduce that, to leading order, the g_n 's must behave as modular forms of $\Gamma_1(3)$ with weight $3n + 1$. Indeed,

$$S'(g_n(\tau, T_\ell)) := g_n\left(\frac{\tau}{1 - 3\tau}, S'(T_\ell)\right) = (1 - 3\tau)^{3n+1}((g_n(\tau, T_\ell) + \dots)). \quad (5.10)$$

The modular forms of $\Gamma_1(3)$ form a ring generated by two elements: $f_1(\tau) = (f_+(\tau))^{1/3}$, which has weight 1, and $f_-(\tau)$, which has weight 3. Here $f_\pm(\tau)$ are the functions defined in (3.20). As we will see in more detail later, the fact that the above equation holds only at leading order and that there are corrections as the ellipses in (5.10) indicate, implies that the g_n 's are actually *quasi*-modular functions of $\Gamma_1(3)$. By this we mean that the g_n 's are polynomials in $f_1(\tau)$, $f_-(\tau)$ and $E_2(\tau)$, where $E_2(\tau)$ is the second Eisenstein series of weight 2 defined by

$$E_2(\tau) = 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^n = 1 - 24q - 72q^2 - 96q^3 + \dots \quad (5.11)$$

with $\sigma_1(n)$ being the sum of divisors of n . E_2 is not a modular form owing to its anomalous behaviour under the $\mathrm{SL}(2, \mathbb{Z})$ transformations:

$$E_2\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^2 E_2(\tau) + \frac{6c}{i\pi}(c\tau + d). \quad (5.12)$$

The modular forms of $\Gamma_1(3)$ can be divided into two classes according to their parity under the S -transformation $\tau \rightarrow -\frac{1}{3\tau}$. Indeed one has (see Appendix B)

$$\begin{aligned} f_1\left(-\frac{1}{3\tau}\right) &= -(\mathrm{i}\sqrt{3}\tau) f_1(\tau) , \\ f_{-}\left(-\frac{1}{3\tau}\right) &= +(\mathrm{i}\sqrt{3}\tau)^3 f_{-}(\tau) , \end{aligned} \quad (5.13)$$

so that we can assign S -parity $(+1)$ to f_{-} , and S -parity (-1) to f_{+} . Since all modular forms of $\Gamma_1(3)$ are generated by these basis elements, the above assignments are enough to specify the S -parity of any modular form of $\Gamma_1(3)$. Furthermore, one can show that the combination

$$\tilde{E}_2(\tau) = E_2(\tau) + f_1^2(\tau) \quad (5.14)$$

has S -parity (-1) since, as we show in Appendix B,

$$\tilde{E}_2\left(-\frac{1}{3\tau}\right) = -(\mathrm{i}\sqrt{3}\tau)^2 \left(\tilde{E}_2(\tau) + \frac{6}{\mathrm{i}\pi\tau}\right) . \quad (5.15)$$

We are now adequately prepared to understand how X transforms under S -duality.

5.1.2 The S transformation

The S -transformation (4.21) implies that the periods in the special vacuum transform as

$$\begin{aligned} a^{\mathrm{D}} &\rightarrow -a , \\ a &\rightarrow -a^{\mathrm{D}} . \end{aligned} \quad (5.16)$$

Therefore, from (5.6) we have

$$S(X) = \frac{X}{\mathrm{i}\sqrt{3}\tau} . \quad (5.17)$$

Acting with S on the right hand side of (5.6) we get

$$S(X) = \frac{\sqrt{3}}{2\pi} \sum_{n=0}^{\infty} (-1)^n \frac{S(g_n(\tau, T_\ell))}{(\mathrm{i}\sqrt{3}\tau a)^{3n+2}} \left(1 + \frac{1}{2\pi\mathrm{i}\tau} \sum_m \frac{g_m}{a^{3m+3}}\right)^{-3n-2} . \quad (5.18)$$

Equating (5.17) and (5.18), at leading order we find that

$$g_n\left(-\frac{1}{3\tau}, S(T_\ell)\right) = (-1)^n \left(\mathrm{i}\sqrt{3}\tau\right)^{3n+1} (g_n(\tau, T_\ell) + \dots) . \quad (5.19)$$

From the behaviour of g_n we can now infer that, to leading order, the g_n 's are modular forms of $\Gamma_1(3)$ with parity $(-1)^n$. The sub-leading terms will modify this to the statement that the g_n 's are *quasi*-modular forms of $\Gamma_1(3)$, with definite parity.

Let us now give some details. In the simplest case of g_0 , there are no subleading corrections to the S and S' transformations. Our previous argument tells us that g_0 is a modular form of $\Gamma_1(3)$ with weight 1 and S -parity $+1$. As mentioned above, there is a unique modular form of weight 1, namely

$$f_1(\tau) = (f_+(\tau))^{\frac{1}{3}} = 1 + 6q + 6q^2 + 6q^3 + \dots . \quad (5.20)$$

While this perfectly agrees with the q -expansion in (5.5), f_1 has S -parity -1 . Therefore, in order for the coefficients to match, the T_ℓ 's must also transform under the S -transformation according to

$$S(T_3) = -T_3 \quad \text{and} \quad S(T_6) = T_6 . \quad (5.21)$$

This completes the identification of g_0 as a modular form of weight 1 and S -parity $+1$, namely

$$g_0(\tau, T_\ell) = -\frac{T_3}{6} f_1(\tau) . \quad (5.22)$$

One could repeat this analysis for the other g_n 's. However, it is more convenient and efficient to exploit a recursion relation satisfied by these coefficients, to which we now turn.

5.2 Recursion relation

Before stating the general result, let us work out the explicit dual transformation properties of the lower g_n 's. Since the behaviour under S and S' are very similar, we will only exhibit the transformation rules under S which are a bit simpler to discuss. We begin by comparing the coefficient of the term proportional to $1/a^5$ in the two expressions (5.17) and (5.18) for $S(X)$. From the first one we have

$$S(X) \Big|_{1/a^5} = \frac{1}{2\pi i \tau} g_1 , \quad (5.23)$$

while from the second we get

$$S(X) \Big|_{1/a^5} = -\frac{\sqrt{3}}{2\pi} \left[\frac{S(g_1)}{(i\sqrt{3}\tau)^5} + \frac{S(g_0)}{(i\sqrt{3}\tau)^2} \frac{g_0}{\pi i \tau} \right] . \quad (5.24)$$

Using the fact that $S(g_0) = (i\sqrt{3}\tau) g_0$, as one can see from (5.22), the above two expressions imply

$$S(g_1) = -(i\sqrt{3}\tau)^4 \left(g_1 + \frac{g_0^2}{\pi i \tau} \right) . \quad (5.25)$$

The presence of the g_0^2 term clearly shows that g_1 cannot be simply a modular form; in fact to compensate for its presence, g_1 must be a quasi-modular form, as we have already anticipated. The only quasi-modular form is the second Eisenstein series E_2 or its modified version \tilde{E}_2 defined in (5.14). To stress the fact that \tilde{E}_2 should appear in the expression for g_1 , we temporarily modify the notation and, in place of $g_1(\tau, T_\ell)$, we write

$$g_1(\tilde{E}_2, T_3, T_6) . \quad (5.26)$$

We instead continue to leave implicit the dependence on the other modular forms to avoid clutter in the formulas. Then, using the known S -transformation of \tilde{E}_2 (see (5.15)), and demanding that g_1 has modular weight 4 and S -parity (-1) , we have

$$\begin{aligned} S(g_1) &:= g_1(S(\tilde{E}_2), S(T_3), S(T_6)) = g_1\left(\tilde{E}_2(-\tfrac{1}{3\tau}), -T_3, T_6\right) \\ &= -(i\sqrt{3}\tau)^4 g_1\left(\tilde{E}_2 + \frac{6}{\pi i \tau}, -T_3, T_6\right) \\ &= -(i\sqrt{3}\tau)^4 \left[g_1(\tilde{E}_2, -T_3, T_6) + \frac{6}{\pi i \tau} \frac{\partial}{\partial \tilde{E}_2} g_1(\tilde{E}_2, -T_3, T_6) + \dots \right] \end{aligned} \quad (5.27)$$

where we have used (5.15) and (5.21) and the dots stand for terms of higher order in $1/\tau$ which are proportional to higher \tilde{E}_2 -derivatives. Comparing with (5.25), we deduce that g_1 must be an even function of T_3 and that, in order to match the $1/\tau$ terms, the following relation must hold

$$\frac{\partial g_1}{\partial \tilde{E}_2} = \frac{1}{6} g_0^2 . \quad (5.28)$$

Integrating with respect to \tilde{E}_2 , we find that

$$g_1 = \frac{T_3^2}{216} f_1^2 \tilde{E}_2 + g_1^{(0)} \quad (5.29)$$

where $g_1^{(0)}$ is a modular form of $\Gamma_1(3)$ with weight 4 and S -parity -1 . As shown in Tab. 2 of Appendix B, there is only one such modular form, namely $f_1 f_-$, and thus $g_1^{(0)}$ must be proportional to it. The proportionality coefficient can be fixed by comparing with the perturbative value of the $1/a^5$ term of X , which is $T_6/30$ as one can deduce from (5.3). The final result for g_1 is then

$$g_1 = \frac{T_3^2}{216} (f_1^2 \tilde{E}_2 - 2 f_1 f_-) + \frac{T_6}{30} f_1 f_- . \quad (5.30)$$

Notice that this is an even function of T_3 as it should be, and that it is linear in \tilde{E}_2 . Thus in (5.27) there are no higher order terms in $1/\tau$, in perfect agreement with (5.25).

By expanding the modular forms in powers of q we can obtain the multi-instanton contributions to g_1 and check that they perfectly agree with the explicit results reported in (5.5) and obtained with the equivariant localization methods. The fact that by using the differential equation (5.28) and *only* the 1-loop result it is possible to get the entire tail of instanton corrections and the fact that these are in complete agreement with the explicit multi-instanton calculations are a very strong and highly non-trivial consistency check of our analysis.

The process we have described can be repeated order by order in n for the higher coefficients g_n , and the final result can be compactly stated in the form of a recursion relation:

$$\frac{\partial g_n}{\partial \tilde{E}_2} = \frac{(3n+1)}{24} \sum_{\ell < n} g_\ell g_{n-\ell-1} . \quad (5.31)$$

Using this relation, one can determine the quasi-modular parts of g_n . The remaining terms are fixed in terms of modular functions of $\Gamma_1(3)$ with definite S -parity by comparing the first few coefficients of the q -expansion. For example, at the next order we find

$$g_2 = -\frac{T_3^3}{1296} \left(\frac{7 f_1^3 \tilde{E}_2^2}{24} - \frac{7 f_1^2 f_- \tilde{E}_2}{6} + \frac{11 f_1^7}{18} - \frac{7 f_1 f_-^2}{9} \right) - \frac{T_3 T_6}{144} \left(\frac{7 f_1^2 f_- \tilde{E}_2}{15} + \frac{3 f_1^7}{10} - \frac{7 f_1 f_-^2}{30} \right) . \quad (5.32)$$

To fix the coefficients of the modular forms f_1^7 and $f_1 f_-^2$ we have matched the 1-loop and the 1-instanton terms, while the higher instanton corrections are predicted by the q -expansion of the right hand side. Again the perfect agreement with the explicit results from localization is a highly non-trivial consistency check.

Using this method we also computed g_3 and g_4 . Their expressions are given in Appendix C.

6. Summary and discussion

In this work we have studied the $\mathcal{N} = 2$ supersymmetric conformal $SU(3)$ gauge theory with six fundamental flavours restricted to a \mathbb{Z}_3 -symmetric locus of its Coulomb branch. The main results are based on the calculation of the prepotential, the dual periods and the period matrix in this special vacuum. These include both one-loop and instanton contributions (up to instanton number 4) to these observables, for general mass configurations.

For a restricted \mathbb{Z}_3 -invariant mass configuration, the period matrix was shown to be proportional to the Cartan matrix of $SU(3)$:

$$\Omega = \tau(q_0) \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} . \quad (6.1)$$

The proportionality constant, in the massless limit, defines an effective τ -parameter, and we showed that the S-duality group acts faithfully on it. We further obtained the S-duality action on the relevant period integrals (a, a^D) in the special vacuum. Combining this with the general form of the dual periods in the (large- a) semi-classical expansion,

$$a^D = i\sqrt{3}\tau a + \frac{\sqrt{3}}{2\pi} \sum_n \frac{g_n(\tau, T_k)}{a^{3n+2}} , \quad (6.2)$$

we showed that the transformation of τ and the period integrals (a, a_D) under the S-duality group constrains the coefficients $g_n(\tau)$ to be quasi-modular forms of $\Gamma_1(3)$. Effectively, this amounts to resumming all instanton contributions at a given order in the large- a expansion. The quasi-modular part of g_n , namely the part that depends on the second Eisenstein series $E_2(\tau)$ or its extension $\tilde{E}_2(\tau)$ defined in (5.14), is completely determined by means of the following recursion relation:

$$\frac{\partial g_n}{\partial \tilde{E}_2} = \frac{(3n+1)}{24} \sum_{\ell < n} g_\ell g_{n-\ell-1} . \quad (6.3)$$

This determines g_n up to modular functions of $\Gamma_1(3)$ with a given weight $3n+1$.

Of particular note in our derivation of quasi-modularity is the role played by the full S-duality group and not just the subgroup $\Gamma_1(3)$. There are numerous modular forms of $\Gamma_1(3)$ for a given weight (see Tab. 2 in Appendix B) and naively, one would have expected that the knowledge of the instanton expansion to a very high order were necessary to fix the modular part completely. However, the constraint from the S -transformation only allows for those modular forms of $\Gamma_1(3)$ that have fixed S -parity to appear in the expansion of the dual periods. Taking this into account, it proves sufficient to use only the perturbative and 1-instanton contribution to completely determine the first few g_n coefficients. Remarkably, the higher order instanton calculations are completely consistent with the resulting expansion in terms of quasi-modular functions.

It would be interesting to extend our analysis away from the special vacuum. At a generic point in the moduli space, the period matrix is no longer proportional to the Cartan matrix. Given that the S-duality group acts as a subgroup of $Sp(4, \mathbb{Z})$, we would expect

that duality would constrain calculable quantities to have an expansion in terms of genus-2 theta functions. Similarly, turning on the Ω -deformation parameters would also take us away from the special vacuum and it would be interesting to see whether there is any sort of quasi-modular behaviour of the prepotential or dual periods in such cases.

Once our analysis is extended to a generic point in the Coulomb moduli space, it would be interesting to explore whether our results could be useful in studying the infinite coupling point in the moduli space of couplings [28]. Since the exact dependence on the coupling constant is known, one could take the coupling to be close to the infinite coupling limit and directly derive properties of the dual theories. This could give us useful information about non-Lagrangian theories, such as the superconformal theories with E_n symmetry [29, 30], that would be difficult to obtain otherwise.

Finally, the S-duality group of superconformal quivers has been studied in great detail in [31]. Furthermore the AGT-W correspondence relates the partition function of the Ω -deformed gauge theory to chiral conformal blocks in two dimensional Toda CFT [32, 33]. In the $SU(2)$ case, recursion relations that encapsulate the quasi-modular structure of the prepotential have been obtained in [34, 35, 36] by studying null-vector decoupling equations in Liouville theory. It would be interesting to see if one can recover the recursion relations derived in this paper by studying similar equations in Toda theory.

Acknowledgments

We thank Carlo Angelantonj, Soumyadeep Bhattacharya, Francesco Fucito, Laurent Gallot, Hossein Parsa Ghorbani, Dileep Jatkar and Renjan John for several discussions.

The work of M.B., M.F. and A.L. is partially supported by the Compagnia di San Paolo contract “MAST: Modern Applications of String Theory” TO-Call3-2012-0088.

A. Observables in the special vacuum

In this appendix we collect the explicit expressions for some of the observables of the $SU(3)$ theory in the special vacuum obtained using equivariant localization. All observables are first calculated at a generic point in the $U(3)$ moduli space after which the special vacuum constraint (2.17) is imposed.

The prepotential coefficients

In the special vacuum the prepotential takes the form

$$F \Big|_{\text{s.v.}} = -\hat{h}_0 \log \left(\frac{a^3}{\Lambda^3} \right) + \sum_{n=1}^{\infty} \frac{\hat{h}_n(q_0, T_\ell)}{n a^{3n}}. \quad (\text{A.1})$$

Up to four instantons, the first few coefficients $\widehat{h}_n(q_0, T_\ell)$ are

$$\widehat{h}_0 = -\frac{T_2}{2} , \quad (\text{A.2a})$$

$$\begin{aligned} \widehat{h}_1 = & \frac{T_5}{131220} (6561 + 8748q_0 + 3402q_0^2 + 2028q_0^3 + 1412q_0^4 + \dots) \\ & - \frac{T_2 T_3}{39366} (2187q_0 + 972q_0^2 + 603q_0^3 + 428q_0^4 + \dots) , \end{aligned} \quad (\text{A.2b})$$

$$\begin{aligned} \widehat{h}_2 = & -\frac{T_2^4}{5878656} (2187 - 2268q_0^2 - 2464q_0^3 - 2156q_0^4 + \dots) \\ & + \frac{T_2^2 T_4}{1469664} (6561 - 6804q_0^2 - 7392q_0^3 - 6692q_0^4 + \dots) \\ & - \frac{T_4^2}{1469664} (6561 + 6804q_0^2 + 4704q_0^3 + 3500q_0^4 + \dots) \\ & - \frac{T_3 T_5}{229635} (2187 - 567q_0^2 - 742q_0^3 - 700q_0^4 + \dots) \\ & + \frac{T_2 T_3^2}{551124} (2187 - 3402q_0^2 - 3472q_0^3 - 3080q_0^4 + \dots) \\ & - \frac{T_2 T_6}{183708} (2187 - 2268q_0^2 - 2128q_0^3 - 1820q_0^4 + \dots) . \end{aligned} \quad (\text{A.2c})$$

As they stand these expressions are not very illuminating. Indeed, it is more useful to replace the instanton counting parameter q_0 and rewrite them in terms of the effective coupling q using the UV/IR relation (2.28). Denoting the new coefficients obtained in this way as $h(q, T_\ell)$, we have

$$h_0 = -\frac{T_2}{2} , \quad (\text{A.3a})$$

$$\begin{aligned} h_1 = & \frac{T_5}{20} (1 - 36q - 54q^2 - 252q^3 - 468q^4 + \dots) \\ & + \frac{3T_2 T_3}{2} (q + 3q^3 + 4q^4 + \dots) , \end{aligned} \quad (\text{A.3b})$$

$$\begin{aligned} h_2 = & -\frac{T_2^4}{2688} (1 - 756q^2 + 4032q^3 + 29484q^4 + \dots) \\ & + \frac{T_2^2 T_4}{224} (1 - 756q^2 + 4032q^3 + 11340q^4 + \dots) \\ & - \frac{T_4^2}{224} (1 + 756q^2 + 4032q^3 + 20412q^4 + \dots) \\ & - \frac{T_3 T_5}{105} (1 - 189q^2 + 2142q^3 + 9072q^4 + \dots) \\ & + \frac{T_2 T_3^2}{252} (1 - 1134q^2 + 4032q^3 + 9072q^4 + \dots) \\ & - \frac{T_6 T_2}{84} (1 - 756q^2 + 1008q^3 + 2268q^4 + \dots) . \end{aligned} \quad (\text{A.3c})$$

We observe that for the mass configuration (2.23) for which only the Casimirs T_3 and T_6 are non-zero, all coefficients \widehat{h}_n (or h_n) vanish.

The dual period coefficients

We present the results for the dual period $a^D := \frac{1}{2\pi i} \frac{\partial F}{\partial a_2} \Big|_{\text{s.v.}}$ in the special vacuum. This is initially obtained as an expansion in the microscopic coupling q_0 . As for the prepotential, the result takes a much simpler form when reexpressed in terms of the effective coupling q :

$$a^D = i\sqrt{3}\tau a + \frac{\sqrt{3}}{2\pi} \sum_{n=0}^{\infty} \left[\frac{h_n(q, T_\ell)}{a^{3n+1}} + \frac{g_n(q, T_\ell)}{a^{3n+2}} \right] \quad (\text{A.4})$$

where the functions $h_n(q, T_\ell)$ are the same as in (A.3) while the first few functions $g_n(q, T_\ell)$ up to 4 instantons are given by

$$g_0 = -\frac{T_3}{6} (1 + 6q + 6q^3 + 6q^4 + \dots) , \quad (\text{A.5a})$$

$$\begin{aligned} g_1 = & -\frac{T_2^3}{8} (q + 27q^3 - 224q^4 + \dots) \\ & + \left(\frac{T_3^2}{3} + \frac{3T_2T_4}{4} \right) (q + 6q^2 + 3q^3 + 28q^4 + \dots) \\ & + \frac{T_6}{30} (1 - 30q - 270q^2 - 570q^3 - 2190q^4 + \dots) , \end{aligned} \quad (\text{A.5b})$$

$$\begin{aligned} g_2 = & \frac{T_2^3T_3}{576} (1 - 468q^2 - 5952q^3 + 4914q^4 + \dots) \\ & + \frac{T_3^3}{1296} (1 - 756q^2 - 336q^3 + 35154q^4 + \dots) \\ & + \frac{3T_2T_3T_4}{4} (q^2 + 26q^3 + 186q^4 + \dots) \\ & - \frac{T_2^2T_5}{320} (1 - 396q^2 - 6528q^3 + 1026q^4 + \dots) \\ & - \frac{T_4T_5}{160} (1 + 396q^2 + 5856q^3 + 35586q^4 + \dots) \\ & - \frac{T_3T_6}{144} (1 - 252q^2 + 672q^3 + 20034q^4 + \dots) . \end{aligned} \quad (\text{A.5c})$$

When we set $T_2 = T_4 = T_5 = 0$, we obtain the result quoted in (5.5).

The period matrix coefficients

In the special vacuum the period matrix can be put in the following form

$$\Omega = \tau \mathcal{C} - \frac{1}{2\pi i} \sum_{n=0}^{\infty} \left[\left(\frac{3n+1}{a^{3n+2}} h_n(q, T_\ell) \right) \mathcal{B} + \left(\frac{3n+2}{a^{3n+3}} g_n(q, T_\ell) \right) \mathcal{C} + \left(\frac{3n+3}{a^{3n+4}} k_n(q, T_\ell) \right) \mathcal{B}^\dagger \right] \quad (\text{A.6})$$

where \mathcal{C} is the Cartan matrix of $\text{SU}(3)$, and the matrices \mathcal{B} and \mathcal{B}^\dagger are defined in (2.21). One could, of course, write the coefficients in terms of the bare coupling constant q_0 , leading to hatted coefficient-functions. But we choose to present the period matrix in terms of the effective q parameter since the results take a simple form. In (A.6), the functions $h_n(q, T_\ell)$

and $g_n(q, T_\ell)$ are the same ones obtained above while the $k_n(q, T_\ell)$ are given by

$$k_0 = \frac{T_2^2}{2} (q - 6q^2 + 57q^3 - 452q^4 + \dots) - \frac{T_4}{12} (1 + 12q + 36q^2 + 12q^3 + 84q^4 + \dots) , \quad (\text{A.7a})$$

$$k_1 = -\frac{T_2^2 T_3}{144} (1 - 162q^2 - 2400q^3 + 1512q^4) + \frac{T_3 T_4}{72} (1 - 162q^2 - 1248q^3 - 4536q^4) + \frac{T_2 T_5}{60} (1 - 1788q^3 + 2916q^4 + \dots) , \quad (\text{A.7b})$$

$$k_2 = -\frac{T_2^5}{8640} (1 - 540q^2 - 5100q^3 - 79380q^4 + \dots) + \frac{T_2^3 T_4}{864} (1 - 432q^2 - 8592q^3 - 39096q^4 + \dots) - \frac{T_3^2 T_4}{648} (1 - 216q^2 - 1752q^3 - 1512q^4) - \frac{T_5^2}{450} (1 + 378q^2 + 6492q^3 + 74196q^4) + \frac{T_2 T_4^2}{4} (3q^2 + 67q^3 + 411q^4 + \dots) - \frac{T_2 T_3 T_5}{270} (1 - 378q^2 - 5370q^3 - 27756q^4 + \dots) + \frac{T_2^2 T_3^2}{432} (1 - 324q^2 - 3528q^3 - 11376q^4 + \dots) - \frac{T_2^2 T_6}{432} (1 - 216q^2 - 6720q^3 - 39744q^4) - \frac{T_4 T_6}{216} (1 + 216q^2 + 6384q^3 + 60912q^4) . \quad (\text{A.7c})$$

B. Modular forms of $\Gamma_1(3)$

In this appendix we collect some useful formulas on the modular and quasi-modular forms used in the main text.

We first define $f_\pm(\tau)$ to be

$$f_\pm(\tau) = \left(\frac{\eta^3(\tau)}{\eta(3\tau)} \right)^3 \pm \left(\frac{3\eta^3(3\tau)}{\eta(\tau)} \right)^3 , \quad (\text{B.1})$$

where the Dedekind η -function is

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n) . \quad (\text{B.2})$$

According to [22, 23], the modular functions of $\Gamma_1(3)$ (the congruence subgroup of $\text{Sl}(2, \mathbb{Z})$ defined in (4.25)) form a ring generated by the two basis elements: $\{f_1(\tau), f_-(\tau)\}$, where

$$f_1(\tau) = (f_+(\tau))^{\frac{1}{3}} . \quad (\text{B.3})$$

These functions have the following Fourier expansions for small $q = e^{2\pi i\tau}$:

$$\begin{aligned} f_1 &= 1 + 6q + 6q^3 + 6q^4 + \dots , \\ f_- &= 1 - 36q - 54q^2 - 252q^3 - 468q^4 + \dots . \end{aligned} \quad (\text{B.4})$$

Note that f_1 is a modular form of $\Gamma_1(3)$ with weight 1, while f_- is a modular form of $\Gamma_1(3)$ with weight 3. This means is that under the S' -transformation

$$S' : \tau \rightarrow \frac{\tau}{1 - 3\tau} , \quad (\text{B.5})$$

these functions behave as follows

$$\begin{aligned} f_1\left(\frac{\tau}{1 - 3\tau}\right) &= (1 - 3\tau) f_1(\tau) , \\ f_-\left(\frac{\tau}{1 - 3\tau}\right) &= (1 - 3\tau)^3 f_-(\tau) . \end{aligned} \quad (\text{B.6})$$

These modular forms have nice transformation properties also under the S -transformation

$$S : \tau \rightarrow -\frac{1}{3\tau} , \quad (\text{B.7})$$

which lies outside $\Gamma_1(3)$. To show this, we can exploit the known transformation of the Dedekind η -function

$$\eta\left(-\frac{1}{\tau}\right) = \sqrt{-i\tau} \eta(\tau) , \quad (\text{B.8})$$

and check that the functions f_{\pm} transform as

$$f_{\pm}\left(-\frac{1}{3\tau}\right) = \mp(i\sqrt{3}\tau)^3 f_{\pm}(\tau) . \quad (\text{B.9})$$

From this we get the S -transformation of f_1 which clearly is only determined up to a \mathbb{Z}_3 phase factor that we choose it to be 1. Thus,

$$f_1\left(-\frac{1}{3\tau}\right) = - (i\sqrt{3}\tau) f_1(\tau) . \quad (\text{B.10})$$

Basis of quasi-modular forms

We now list the basis of quasi-modular forms of $\Gamma_1(3)$ which have been used in the main text. In addition to the weight under the $\Gamma_1(3)$ transformations, we have a \mathbb{Z}_2 charge, which we have call S -parity, determined by the properties under the S -transformation. For instance, from the transformation properties (B.10) and (B.9) we assign f_1 an S -parity (-1) and f_- an S -parity $(+1)$. In addition to these modular forms, we also have the quasi-modular

form $E_2(\tau)$. However, E_2 by itself, does not transform well under the S -transformation; indeed

$$E_2\left(-\frac{1}{3\tau}\right) = (3\tau)^2 E_2(3\tau) + \frac{6}{i\pi}(3\tau) . \quad (\text{B.11})$$

On the other hand one can prove the following triplication formula

$$E_2(3\tau) = \frac{1}{3}E_2(\tau) + \frac{2}{3}f_1^2(\tau) \quad (\text{B.12})$$

which shows that E_2 mixes with f_1^2 under the S -transformation. Using this fact and the S -transformation property of f_1 given in (B.10), it is possible to check that the following combination

$$\tilde{E}_2(\tau) = E_2(\tau) + f_1^2(\tau) \quad (\text{B.13})$$

transforms into itself, save for an anomalous term:

$$\tilde{E}_2\left(-\frac{1}{3\tau}\right) = -(i\sqrt{3}\tau)^2 \tilde{E}_2(\tau) + \frac{18}{i\pi}\tau . \quad (\text{B.14})$$

Thus, we can assign the quasi-modular form \tilde{E}_2 an S -parity equal to -1 .

In Tab. 2 we list a basis of the modular forms of $\Gamma_1(3)$ up to weight 10, distinguishing them according to their S -parity.

Weight	Modular forms of $\Gamma_1(3)$	S -parity
1	f_1	-1
2	f_1^2	$+1$
3	f_1^3	-1
	f_-	$+1$
4	$f_1 f_-$	-1
	f_1^4	$+1$
5	f_1^5	-1
	$f_1^2 f_-$	$+1$
6	$f_1^3 f_-$	-1
	f_1^6, f_-^2	$+1$
7	$f_1^7, f_1 f_-^2$	-1
	$f_1^4 f_-$	$+1$
8	$f_1^5 f_-$	-1
	$f_1^8, f_1^2 f_-^2$	$+1$
9	$f_1^9, f_1^3 f_-^2$	-1
	$f_1^6 f_-, f_-^3$	$+1$
10	$f_1^7 f_-, f_1 f_-^3$	-1
	$f_1^{10}, f_1^4 f_-^2$	$+1$

Table 2: A basis of modular forms of $\Gamma_1(3)$ up to weight 10, classified according to their S -parity.

C. The coefficients g_3 and g_4

Here we report the expression of the coefficients g_3 and g_4 in the expansion of the dual period, written in terms of the quasi-modular functions of $\Gamma_1(3)$, which we obtained by solving the recursion relation (5.31).

$$\begin{aligned}
g_3 = & -\frac{T_3^4}{7128} \left(\frac{605f_1^8\tilde{E}_2}{1296} - \frac{55f_1^4\tilde{E}_2^3}{576} - \frac{323f_1^7f_-}{648} + \frac{55f_1^3\tilde{E}_2^2f_-}{96} - \frac{275f_1^2\tilde{E}_2f_-^2}{324} + \frac{119f_1f_-^3}{162} \right) \\
& + \frac{T_3^2T_6}{1188} \left(\frac{11f_1^8\tilde{E}_2}{32} - \frac{73f_1^7f_-}{144} + \frac{11f_1^3\tilde{E}_2^2f_-}{32} - \frac{55f_1^2\tilde{E}_2f_-^2}{96} + \frac{85f_1f_-^3}{144} \right) \\
& + \frac{T_6^2}{792} \left(\frac{7f_1^7f_-}{10} + \frac{11f_1^2\tilde{E}_2f_-^2}{30} - \frac{13f_1f_-^3}{30} \right), \tag{C.1a}
\end{aligned}$$

$$\begin{aligned}
g_4 = & -\frac{T_3^5}{68040} \left(\frac{5435f_1^{13}}{21384} - \frac{5005f_1^9\tilde{E}_2^2}{10368} + \frac{5005f_1^5\tilde{E}_2^4}{82944} + \frac{16835f_1^8\tilde{E}_2f_-}{14256} - \frac{5005f_1^4\tilde{E}_2^3f_-}{10368} \right. \\
& \quad \left. - \frac{91351f_1^7f_-^2}{85536} + \frac{5915f_1^3\tilde{E}_2^2f_-^2}{5184} - \frac{15925f_1^2\tilde{E}_2f_-^3}{9504} + \frac{182455f_1f_-^4}{171072} \right) \\
& + \frac{T_3^3T_6}{108} \left(\frac{13f_1^{13}}{3960} - \frac{13f_1^9\tilde{E}_2^2}{3840} + \frac{10621f_1^8\tilde{E}_2f_-}{855360} - \frac{143f_1^4\tilde{E}_2^3f_-}{51840} - \frac{7631f_1^7f_-^2}{855360} \right. \\
& \quad \left. + \frac{299f_1^3\tilde{E}_2^2f_-^2}{34560} - \frac{11791f_1^2\tilde{E}_2f_-^3}{855360} + \frac{8021f_1f_-^4}{855360} \right) \\
& - \frac{T_3T_6^2}{1512} \left(\frac{237f_1^{13}}{2200} + \frac{9373f_1^8\tilde{E}_2f_-}{26400} - \frac{5161f_1^7f_-^2}{39600} + \frac{91}{600}f_1^3\tilde{E}_2^2f_-^2 \right. \\
& \quad \left. - \frac{2093f_1^2\tilde{E}_2f_-^3}{8800} + \frac{7189f_1f_-^4}{39600} \right). \tag{C.1b}
\end{aligned}$$

As is evident from the above expressions and in the earlier discussion on recursion relations, the quasi-modular piece of these expressions – terms with \tilde{E}_2 dependence – are completely fixed by the recursion relation. The remaining pieces are decided by considerations of weight and S -parity. Thus, to fix g_3 one needs to use the perturbative and 1-instanton contributions, while to fix g_4 , one needs to take into account 2-instanton corrections as well.

By expanding the quasi-modular forms in powers of q one can check that the expressions agree with the instanton corrections up to $k = 4$ given in (5.5d) and (5.5e).

References

- [1] J. Teschner, *Exact results on $N=2$ supersymmetric gauge theories*, [arXiv:1412.7145 \[hep-th\]](#).
- [2] M.-x. Huang, A.-K. Kashani-Poor, and A. Klemm, *The Ω deformed B -model for rigid $\mathcal{N} = 2$ theories*, *Annales Henri Poincaré* **14** (2013) 425–497, [arXiv:1109.5728 \[hep-th\]](#).

- [3] M. Billo, M. Frau, L. Gallot, A. Lerda, and I. Pesando, *Deformed $N=2$ theories, generalized recursion relations and S -duality*, JHEP **1304** (2013) 039, [arXiv:1302.0686 \[hep-th\]](#).
- [4] M. Billo, M. Frau, L. Gallot, A. Lerda, and I. Pesando, *Modular anomaly equation, heat kernel and S -duality in $N = 2$ theories*, JHEP **1311** (2013) 123, [arXiv:1307.6648 \[hep-th\]](#).
- [5] M.-x. Huang, *Modular anomaly from holomorphic anomaly in mass deformed $N=2$ superconformal field theories*, Phys. Rev. D **87** (2013) 105010, [arXiv:1302.6095 \[hep-th\]](#).
- [6] M. Billo, M. Frau, F. Fucito, A. Lerda, J. Morales, R. Poghossyan, and D. Ricci-Pacifici, *Modular anomaly equations in $\mathcal{N} = 2^*$ theories and their large- N limit*, JHEP **1410** (2014) 131, [arXiv:1406.7255 \[hep-th\]](#).
- [7] M. Billo, M. Frau, F. Fucito, A. Lerda and J. F. Morales, *S -duality and the prepotential in $N=2^*$ theories (I): the ADE algebras*, [arXiv:1507.07709 \[hep-th\]](#).
- [8] M. Billo, M. Frau, F. Fucito, A. Lerda and J. F. Morales, *S -duality and the prepotential in $N=2^*$ theories (II): the non-simply laced algebras*, [arXiv:1507.08027 \[hep-th\]](#).
- [9] N. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, Adv. Theor. Math. Phys. **7** (2004) 831–864, [arXiv:hep-th/0206161](#).
- [10] N. Nekrasov and A. Okounkov, *Seiberg-Witten theory and random partitions*, [arXiv:hep-th/0306238](#).
- [11] U. Bruzzo, F. Fucito, J. F. Morales, and A. Tanzini, *Multi-instanton calculus and equivariant cohomology*, JHEP **05** (2003) 054, [arXiv:hep-th/0211108](#).
- [12] F. Fucito, J. F. Morales, and R. Poghossian, *Multi instanton calculus on ALE spaces*, Nucl.Phys. **B703** (2004) 518–536, [arXiv:hep-th/0406243 \[hep-th\]](#).
- [13] N. Seiberg and E. Witten, *Electric - magnetic duality, monopole condensation, and confinement in $N=2$ supersymmetric Yang-Mills theory*, Nucl.Phys. **B426** (1994) 19–52, [arXiv:hep-th/9407087 \[hep-th\]](#).
- [14] N. Seiberg and E. Witten, *Monopoles, duality and chiral symmetry breaking in $N=2$ supersymmetric QCD*, Nucl. Phys. **B431** (1994) 484–550, [arXiv:hep-th/9408099](#).
- [15] E. Witten, *Solutions of four-dimensional field theories via M theory*, Nucl.Phys. **B500** (1997) 3–42, [arXiv:hep-th/9703166 \[hep-th\]](#).
- [16] A. Hanany and Y. Oz, *On the quantum moduli space of vacua of $N=2$ supersymmetric $SU(N(c))$ gauge theories*, Nucl.Phys. **B452** (1995) 283–312, [arXiv:hep-th/9505075 \[hep-th\]](#).
- [17] P. C. Argyres, M. R. Plesser, and A. D. Shapere, *The Coulomb phase of $N=2$ supersymmetric QCD*, Phys.Rev.Lett. **75** (1995) 1699–1702, [arXiv:hep-th/9505100 \[hep-th\]](#).
- [18] J. A. Minahan and D. Nemeschansky, *Hyperelliptic curves for supersymmetric Yang-Mills*, Nucl. Phys. **B464** (1996) 3–17, [arXiv:hep-th/9507032](#).
- [19] J. A. Minahan and D. Nemeschansky, *$N=2$ superYang-Mills and subgroups of $SL(2, \mathbb{Z})$* , Nucl.Phys. **B468** (1996) 72–84, [arXiv:hep-th/9601059 \[hep-th\]](#).
- [20] P. C. Argyres and S. Pellsand, *Comparing instanton contributions with exact results in $N=2$ supersymmetric scale invariant theories*, JHEP **0003** (2000) 014, [arXiv:hep-th/9911255 \[hep-th\]](#).

- [21] P. C. Argyres and A. Buchel, *The Nonperturbative gauge coupling of $N=2$ supersymmetric theories*, Phys.Lett. **B442** (1998) 180–184, [arXiv:hep-th/9806234](#) [hep-th].
- [22] N. Koblitz, *Introduction to elliptic curves and modular forms, 2nd Edition*,. Springer-Verlag. (1993).
- [23] T. M. Apostol, *Modular functions and Dirichlet series in number theory, 2nd Edition*,. Springer-Verlag. (1990).
- [24] M. Billo, M. Frau, F. Fucito, L. Giaccone, A. Lerda, J. F. Morales, and D. Ricci-Pacifici, *Non-perturbative gauge/gravity correspondence in $N=2$ theories*, JHEP **1208** (2012) 166, [arXiv:1206.3914](#) [hep-th].
- [25] J. A. Minahan, *Duality symmetries for $N=2$ supersymmetric QCD with vanishing Beta functions*, Nucl.Phys. **B537** (1999) 243–259, [arXiv:hep-th/9806246](#) [hep-th].
- [26] S. Ashok, M. Billo, E. Dell’Aquila, M. Frau, R. John, and A. Lerda, *Non-perturbative studies of $N=2$ conformal quiver gauge theories*, Fortsch.Phys. **63** (2015) 259–293, [arXiv:1502.05581](#) [hep-th].
- [27] D. Gaiotto, G. W. Moore, and A. Neitzke, *Wall-crossing, Hitchin Systems, and the WKB Approximation*, [arXiv:0907.3987](#) [hep-th].
- [28] P. C. Argyres and N. Seiberg, *S -duality in $N=2$ supersymmetric gauge theories*, JHEP **0712** (2007) 088, [arXiv:0711.0054](#) [hep-th].
- [29] J. A. Minahan and D. Nemeschansky, *An $N=2$ superconformal fixed point with $E(6)$ global symmetry*, Nucl. Phys. **B482** (1996) 142–152, [arXiv:hep-th/9608047](#) [hep-th].
- [30] J. A. Minahan and D. Nemeschansky, *Superconformal fixed points with $E(n)$ global symmetry*, Nucl. Phys. **B489** (1997) 24–46, [arXiv:hep-th/9610076](#).
- [31] D. Gaiotto, *$N=2$ dualities*, JHEP **1208** (2012) 034, [arXiv:0904.2715](#) [hep-th].
- [32] L. F. Alday, D. Gaiotto, and Y. Tachikawa, *Liouville correlation functions from four-dimensional gauge theories*, Lett. Math. Phys. **91** (2010) 167–197, [arXiv:0906.3219](#) [hep-th].
- [33] N. Wyllard, *$A(N-1)$ conformal Toda field theory correlation functions from conformal $N = 2$ $SU(N)$ quiver gauge theories*, JHEP **0911** (2009) 002, [arXiv:0907.2189](#) [hep-th].
- [34] A.-K. Kashani-Poor and J. Troost, *The toroidal block and the genus expansion*, JHEP **1303** (2013) 133, [arXiv:1212.0722](#) [hep-th].
- [35] A.-K. Kashani-Poor and J. Troost, *Transformations of spherical blocks*, JHEP **1310** (2013) 009, [arXiv:1305.7408](#) [hep-th].
- [36] A.-K. Kashani-Poor and J. Troost, *Quantum geometry from the toroidal block*, JHEP **1408** (2014) 117, [arXiv:1404.7378](#) [hep-th].